

Oral presentation

Co-infection with *Trypanosoma cruzi* (Chagas' disease agent) decreases HIV-1 transcription in human placenta

Guillermina Dolcini^{*1}, María Elisa Solana², Guadalupe Andreani¹, Ana María Celentano², Ana María Donato³, Alberto Del Río³ and Liliana Martínez Peralta¹

Address: ¹National Reference Centre for AIDS, Microbiology Department, School of Medicine, University of Buenos Aires, Buenos Aires, Argentina,

²Laboratory of Parasitology, Microbiology Department, School of Medicine, University of Buenos Aires, Buenos Aires, Argentina and

³Endocrinology Service, Clinic Biochemistry Department, José de San Martín Hospital, School of Pharmacy and Biochemistry, University of Buenos Aires, Buenos Aires, Argentina

* Corresponding author

from Fourth Dominique Dormont International Conference. Host-Pathogen Interactions in Chronic Infections
Paris, France. 13-15 December 2007

Published: 9 April 2008

Retrovirology 2008, 5(Suppl 1):O13 doi:10.1186/1742-4690-5-S1-O13

This abstract is available from: <http://www.retrovirology.com/content/5/S1/O13>

© 2008 Dolcini et al.; licensee BioMed Central Ltd.

Background

Several factors determine the risk of HIV mother-to-child transmission (MTCT), such as co-infections in placentae from HIV-1 positive mothers with other pathogens [1,2]. One of the most important endemic zoonosis in Latin America is Chagas' disease, caused by the protozoa *Trypanosoma cruzi*. MTCT of *T. cruzi* is today one of the main transmission routes in big cities [3]. The aim of the study was to determine whether *T. cruzi* modifies HIV infection at tissue or cellular level of the placenta.

Material and methods

Simple and double infections were carried out on a placental histoculture system (chorionic villi isolated from term placentae from HIV and Chagas negative mothers) and on choriocarcinoma BeWo cell line. We used trypomastigotes of *T. cruzi* (VD lethal strain, isolated from a child with MTCT), either purified from mice blood or from Vero cell cultures, 24h-supernatants of blood and cellular trypomastigotes, and the viral HIV-1DenvLuc+/VSV-G pseudotype. Viral replication was evaluated by luciferase activity quantification. Tissue viability was evaluated by hCG hormone secretion in histoculture supernatant. Quantification of soluble factors protein secretion and mRNA expression in histocultures were carried out by ELISAs and real time PCR respectively.

Results

Whole trypomastigotes, either from mice blood (mean \pm SD; $-92.13\% \pm 4.85$) or from cell cultures ($-97.33\% \pm 0.58$), in co-infection with viral pseudotype decreased luciferase activity in placental histocultures. Similar results were obtained on BeWo cells. When supernatants from blood trypomastigotes were used on placental histocultures, luciferase activity presented a decrease of $-85.50\% \pm 2.12$, while supernatants from culture trypomastigotes presented lower effects ($-65.50\% \pm 2.12$). Tissue viability was not modified by viral and/or parasite infection. In co-infected histocultures, both protein secretion and mRNA expression of IL-10 were down regulated. Surprisingly, RANTES was increased.

Conclusions

Acute infection with *T. cruzi* and HIV-1 in this placenta *in vitro* system as well as in the trophoblast cell line decreased HIV-1 transcription. *T. cruzi* activity on HIV-1 replication seems to be caused not only by active infection but also by soluble factors shed by the parasite. A balance between proinflammatory and inhibitory cytokines/chimiokines might have a role in this phenomenon. Ongoing experiments are being conducted in order to elucidate the mechanisms involved in the impairment of HIV

replication by *T. cruzi* and their role on MTCT of both pathogens.

References

1. Gallagher M, Malhotra I, Mungai PL, Wamachi AN, Kioko JM, Ouma JH, Muchiri E, King CL: **The effects of maternal helminth and malaria infections on mother-to-child HIV transmission.** *AIDS* 2005, **19**:1849-1855.
2. Ayisi JG, van Eijk AM, Newman RD, ter Kuile FO, Shi YP, Yang C, Kolczak MS, Otieno JA, Misore AO, Kager PA, Lal RB, Steketee RW, Nahlen BL: **Maternal malaria and perinatal HIV transmission, western Kenya.** *Emerg Infect Dis* 2004, **10**:643-652.
3. Carlier Y, Pinto Dias JC, Ostermayer Luquetti A, Hontebeyrie M, Torrico F, Truysens C: **Trypanosomiasis américaine ou maladie de Chagas.** *Maladies infectieuses, Paris* 2002, **8-505-A-20**:1-21. Editions Scientifiques et Médicales Elsevier SAS

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:
http://www.biomedcentral.com/info/publishing_adv.asp

